

Gervais School District#1

Radon Test Results

Fixture Location/Description	Fixture ID#	Test Date	Test Results for Radon
Preschool Room 1	8773371	1/9/2019	<03 ±0.2 pCi/L
Preschool Room 2	8773372	1/9/2019	<03 ±0.2 pCi/L
Maintenance Office	8773376	1/9/2019	<03 ±0.2 pCi/L
Business Office	8773367	1/9/2019	<03 ±0.1 pCi/L
Technology Office	8773368	1/9/2019	<03 ±0.2 pCi/L
Daycare	8773373	1/9/2019	<03 ±0.8 pCi/L
Welcome Center	8773370	1/9/2019	<03 ±0.2 pCi/L
Elementary School 1 st Floor 1	8659778	3/31/2018	<0.3 pCi/L
Elementary School 1 st Floor	8659780	3/31/2018	<0.3 pCi/L
Elementary School 1 st Floor 4	8659781	3/31/2018	<0.3 pCi/L
Elementary School Room 5	8659782	3/31/2018	<0.3 pCi/L
Elementary School 1 st Floor 5a	8659783	3/31/2018	0.8 ±0.6 pCi/L
Elementary School 1 st Floor 6	8659784	3/31/2018	<0.3 pCi/L
Elementary School 1 st Floor	8659785	3/31/2018	????
Elementary School 1 st Floor 8	8659786	3/31/2018	<0.3 pCi/L
Elementary School 1 st Floor 9	8659787	3/31/2018	????
Elementary School 1 st Floor 10	8659788	3/31/2018	<0.3 pCi/L
Elementary School Room 11	8659789	3/31/2018	1.4 ± 0.6 pCi/L
Elementary School 1 st Floor 12	8659790	3/31/2018	????
Elementary School 1 st Floor 16	8659791	3/31/2018	<0.3 pCi/L
Elementary School Room 17	8659792	3/31/2018	<0.3 pCi/L
Elementary School Room 18	8659794	3/31/2018	<0.3 pCi/L
Elementary School Room 19	8659795	3/31/2018	<0.3 pCi/L
Elementary School Room 20	8659796	3/31/2018	<0.3 pCi/L
Elementary School Room 21	8659797	3/31/2018	<0.3 pCi/L
Middle School Room 10	20011946	1/25/2021	<0.3 pCi/L
Middle School Room 15	20011951	1/25/2021	<0.3 pCi/L
Middle School Staff Room	20011955	1/29/2021	<0.3 pCi/L
Middle School Boys Locker Room	20011960	1/25/2021	0.5 pCi/L
Middle School Office	20011961	1/25/2021	<0.3 pCi/L
Middle School Girls Locker Room	20011964	1/26/2021	<0.3 pCi/L
Middle School PE Office	20011965	1/25/2021	<0.3 pCi/L
Middle School Bathroom Gym	20011966	1/25/2021	<0.3 pCi/L
Middle School Room 3	20011967	1/25/2021	<0.3 pCi/L
Middle School Room 11	20011978	1/25/2021	<0.3 pCi/L
Middle School Room 12	20011980	1/25/2021	<0.3 pCi/L

Middle School Room 6	20011983	1/25/2021	<0.3 pCi/L
Middle School Room 14	20011986	1/25/2021	<0.3 pCi/L
Middle School Room 8	20011987	1/25/2021	<0.3 pCi/L
Middle School Room 9	20011988	1/25/2021	<0.3 pCi/L
Middle School Room 2	20011994	1/25/2021	<0.3 pCi/L
Middle School Room 7	20011998	1/25/2021	<0.3 pCi/L
Middle School Room 4	20015764	1/29/2021	<0.3 pCi/L
Middle School Room 1	8959786	1/25/2021	<0.3 pCi/L
Middle School Room 13	20011985	1/29/2021	<0.3 pCi/L
Middle School Room 5	20015728	1/29/2021	<0.3 pCi/L

01/18/19 ACTIVATED CHARCOAL RADON TEST #8773367

Radon Test Result: < 0.3 ±0.1 pCi/L

Test Started 01/09/19 at 2:00 pm
Test Ended 01/15/19 at 9:00 am
Closed house conditions maintained during test.

Location 1st Floor



Air Chek, Inc.
PO Box 2000
Naples, NC 28760

www.radon.com

FEB 7 2019

FEB 7 2019



MELISSA OFFICE
290 1ST ST
GERVAIS, OR 97026-2107

Your Test Result

This result has been rounded to one-tenth (0.1) of a pCi/L (picocurie per liter). This test result reflects the amount of radon measured in this sample AFTER it arrived at our laboratory. All analysis calculations are automatically adjusted to reflect the length of test, the amount of moisture in the sample, temperature, time from the end of test, and the amount of radiation measured. If your test kit was used prior to the Use By date, ALL the testing protocols and instructions were carefully followed, and the data recorded properly on the test packet, then it is reasonable to assume this is an accurate assessment of the average level of the radon this sample was exposed to during the test period.

Health Risks

The primary health risk from long-term exposure to radon is lung cancer. The risk of developing a lung cancer from radon exposure depends both on how much radon is present and how long you are exposed to radon. The higher the radon level or the longer the time of exposure, even if the levels are relatively low, the greater the risk. EPA has set an Action Level for radon at 4 pCi/L; however radon concentrations less than 4 pCi/L still pose some health risks. The Indoor Radon Abatement Act set a goal for indoor radon concentrations to equal the amount of radon found outdoors, which is estimated to be ~ 0.4 pCi/L.

Conducting Follow-up Measurements

USEPA protocol describes two general types of radon measurements: short-term tests conducted from 48 hours up to 90 days, and long-term tests that last from 91 to 365 days. Your first test (initial/screening) should be a short-term 'worst-case' screening to see if there is a potential for high exposure to radon. Screening tests should be conducted under closed-building conditions, in the lowest lived-in area in the house, because the highest concentrations of radon will usually be found in a room closest to the underlying soil. Tests made under these conditions are less likely to miss a house with a potential for high concentrations. On the other hand, if the results of worst-case screening tests are very low, there is a high probability that the average annual concentrations in the house are also low.

(Continued on Back)

INTERPRETING YOUR TEST RESULT

This radon test was provided to you by BLUE RIDGE NET PUBLISHING INC / 540-777-1895. The US EPA action level for indoor radon is 4.0 pCi/L. Test results in this range(0.5 pCi/L or less) are, for all practical purposes, equivalent to the radon levels found in fresh air. However, if you make any structural changes or start to use a lower level of the building more frequently you should test again.

You may be able to obtain additional information about radon related subjects by calling the "Radon Fix-It Line" at 800-644-6999 Monday thru Friday between NOON and 8 pm EST.

Most states have a radon office to assist citizens with general questions about radon and radon reduction techniques. Many states maintain a list of licensed or certified radon testing and mitigation professionals. You can visit www.state-radon.info to find the list of state radon contacts, as well as links to additional radon resources in your area.

01/18/19 ACTIVATED CHARCOAL RADON TEST #8773370

Radon Test Result: < 0.3 ±0.2 pCi/L

Test Started 01/09/19 at 8:00 am

Test Ended 01/15/19 at 9:00 am

Closed house conditions maintained during test.

Location 1st Floor



Air Chek, Inc.

PO Box 2000

Naples, NC 28760

www.radon.com



WELCOME CENTER

345 1ST ST

GERVAIS, OR 97026-

INTERPRETING YOUR TEST RESULT

This radon test was provided to you by BLUE RIDGE NET PUBLISHING INC / 540-777-1895. The US EPA action level for indoor radon is 4.0 pCi/L. Test results in this range(0.5 pCi/L or less) are, for all practical purposes, equivalent to the radon levels found in fresh air. However, if you make any structural changes or start to use a lower level of the building more frequently you should test again.

You may be able to obtain additional information about radon related subjects by calling the "Radon Fix-It Line" at 800-644-6999 Monday thru Friday between NOON and 8 pm EST.

Your Test Result

This result has been rounded to one-tenth (0.1) of a pCi/L (picocurie per liter). This test result reflects the amount of radon measured in this sample AFTER it arrived at our laboratory. All analysis calculations are automatically adjusted to reflect the length of test, the amount of moisture in the sample, temperature, time from the end of test, and the amount of radiation measured. If your test kit was used prior to the Use By date, ALL the testing protocols and instructions were carefully followed, and the data recorded properly on the test packet, then it is reasonable to assume this is an accurate assessment of the average level of the radon this sample was exposed to during the test period.

Health Risks

The primary health risk from long-term exposure to radon is lung cancer. The risk of developing a lung cancer from radon exposure depends both on how much radon is present and how long you are exposed to radon. The higher the radon level or the longer the time of exposure, even if the levels are relatively low, the greater the risk. EPA has set an Action Level for radon at 4 pCi/L; however radon concentrations less than 4 pCi/L still pose some health risks. The Indoor Radon Abatement Act set a goal for indoor radon concentrations to equal the amount of radon found outdoors, which is estimated to be ~ 0.4 pCi/L.

Conducting Follow-up Measurements

USEPA protocol describes two general types of radon measurements: short-term tests conducted from 48 hours up to 90 days, and long-term tests that last from 91 to 365 days. Your first test (initial/screening) should be a short-term 'worst-case' screening to see if there is a potential for high exposure to radon. Screening tests should be conducted under closed-building conditions, in the lowest lived-in area in the house, because the highest concentrations of radon will usually be found in a room closest to the underlying soil. Tests made under these conditions are less likely to miss a house with a potential for high concentrations. On the other hand, if the results of worst-case screening tests are very low, there is a high probability that the average annual concentrations in the house are also low.

(Continued on Back)

Most states have a radon office to assist citizens with general questions about radon and radon reduction techniques. Many states maintain a list of licensed or certified radon testing and mitigation professionals. You can visit www.state-radon.info to find the list of state radon contacts, as well as links to additional radon resources in your area.

01/18/19 ACTIVATED CHARCOAL RADON TEST #8773376

Radon Test Result: < 0.3 ±0.2 pCi/L

Test Started 01/09/19 at 1:00 pm

Test Ended 01/15/19 at 9:00 am

Closed house conditions maintained during test.

Location 1st Floor



Air Chek, Inc.

PO Box 2000

Naples, NC 28760

www.radon.com



TONY'S OFFICE

290 1ST ST

GERVAIS, OR 97026-2107

INTERPRETING YOUR TEST RESULT

This radon test was provided to you by BLUE RIDGE NET PUBLISHING INC / 540-777-1895. The US EPA action level for indoor radon is 4.0 pCi/L. Test results in this range (0.5 pCi/L or less) are, for all practical purposes, equivalent to the radon levels found in fresh air. However, if you make any structural changes or start to use a lower level of the building more frequently you should test again.

You may be able to obtain additional information about radon related subjects by calling the "Radon Fix-It Line" at 800-644-6999 Monday thru Friday between NOON and 8 pm EST.

Your Test Result

This result has been rounded to one-tenth (0.1) of a pCi/L (picocurie per liter). This test result reflects the amount of radon measured in this sample AFTER it arrived at our laboratory. All analysis calculations are automatically adjusted to reflect the length of test, the amount of moisture in the sample, temperature, time from the end of test, and the amount of radiation measured. If your test kit was used prior to the Use By date, ALL the testing protocols and instructions were carefully followed, and the data recorded properly on the test packet, then it is reasonable to assume this is an accurate assessment of the average level of the radon this sample was exposed to during the test period.

Health Risks

The primary health risk from long-term exposure to radon is lung cancer. The risk of developing a lung cancer from radon exposure depends both on how much radon is present and how long you are exposed to radon. The higher the radon level or the longer the time of exposure, even if the levels are relatively low, the greater the risk. EPA has set an Action Level for radon at 4 pCi/L; however radon concentrations less than 4 pCi/L still pose some health risks. The Indoor Radon Abatement Act set a goal for indoor radon concentrations to equal the amount of radon found outdoors, which is estimated to be ~ 0.4 pCi/L.

Conducting Follow-up Measurements

USEPA protocol describes two general types of radon measurements: short-term tests conducted from 48 hours up to 90 days, and long-term tests that last from 91 to 365 days. Your first test (initial/screening) should be a short-term 'worst-case' screening to see if there is a potential for high exposure to radon. Screening tests should be conducted under closed-building conditions, in the lowest lived-in area in the house, because the highest concentrations of radon will usually be found in a room closest to the underlying soil. Tests made under these conditions are less likely to miss a house with a potential for high concentrations. On the other hand, if the results of worst-case screening tests are very low, there is a high probability that the average annual concentrations in the house are also low.

(Continued on Back)

Most states have a radon office to assist citizens with general questions about radon and radon reduction techniques. Many states maintain a list of licensed or certified radon testing and mitigation professionals. You can visit www.state-radon.info to find the list of state radon contacts, as well as links to additional radon resources in your area.

01/18/19 ACTIVATED CHARCOAL RADON TEST #8773368

Radon Test Result: < 0.3 ±0.2 pCi/L

Test Started 01/09/19 at 2:00 pm

Test Ended 01/15/19 at 9:00 am

Closed house conditions maintained during test.

Location 1st Floor



**Air Chek, Inc.
PO Box 2000
Naples, NC 28760**

www.radon.com



**TRACY'S OFFICE
290 1ST ST
GERVAIS, OR 97026-2107**

Your Test Result

This result has been rounded to one-tenth (0.1) of a pCi/L (picocurie per liter). This test result reflects the amount of radon measured in this sample AFTER it arrived at our laboratory. All analysis calculations are automatically adjusted to reflect the length of test, the amount of moisture in the sample, temperature, time from the end of test, and the amount of radiation measured. If your test kit was used prior to the Use By date, ALL the testing protocols and instructions were carefully followed, and the data recorded properly on the test packet, then it is reasonable to assume this is an accurate assessment of the average level of the radon this sample was exposed to during the test period.

INTERPRETING YOUR TEST RESULT

This radon test was provided to you by BLUE RIDGE NET PUBLISHING INC / 540-777-1895. The US EPA action level for indoor radon is 4.0 pCi/L. Test results in this range (0.5 pCi/L or less) are, for all practical purposes, equivalent to the radon levels found in fresh air. However, if you make any structural changes or start to use a lower level of the building more frequently you should test again.

You may be able to obtain additional information about radon related subjects by calling the "Radon Fix-It Line" at 800-644-6999 Monday thru Friday between NOON and 8 pm EST.

Health Risks

The primary health risk from long-term exposure to radon is lung cancer. The risk of developing a lung cancer from radon exposure depends both on how much radon is present and how long you are exposed to radon. The higher the radon level or the longer the time of exposure, even if the levels are relatively low, the greater the risk. EPA has set an Action Level for radon at 4 pCi/L; however radon concentrations less than 4 pCi/L still pose some health risks. The Indoor Radon Abatement Act set a goal for indoor radon concentrations to equal the amount of radon found outdoors, which is estimated to be ~ 0.4 pCi/L.

Conducting Follow-up Measurements

USEPA protocol describes two general types of radon measurements: short-term tests conducted from 48 hours up to 90 days, and long-term tests that last from 91 to 365 days. Your first test (initial/screening) should be a short-term 'worst-case' screening to see if there is a potential for high exposure to radon. Screening tests should be conducted under closed-building conditions, in the lowest lived-in area in the house, because the highest concentrations of radon will usually be found in a room closest to the underlying soil. Tests made under these conditions are less likely to miss a house with a potential for high concentrations. On the other hand, if the results of worst-case screening tests are very low, there is a high probability that the average annual concentrations in the house are also low.

(Continued on Back)

Most states have a radon office to assist citizens with general questions about radon and radon reduction techniques. Many states maintain a list of licensed or certified radon testing and mitigation professionals. You can visit www.state-radon.info to find the list of state radon contacts, as well as links to additional radon resources in your area.

01/18/19 ACTIVATED CHARCOAL RADON TEST #8773371

Radon Test Result: < 0.3 ±0.2 pCi/L

Test Started 01/09/19 at 8:00 am

Test Ended 01/15/19 at 9:00 am

Closed house conditions maintained during test.

Location 1st Floor

FEB 7 2019



Air Chek, Inc.

PO Box 2000

Naples, NC 28760

www.radon.com



PRESCHOOL RM 1

345 1ST ST

GERVAIS, OR 97026

Your Test Result

This result has been rounded to one-tenth (0.1) of a pCi/L (picocurie per liter). This test result reflects the amount of radon measured in this sample AFTER it arrived at our laboratory. All analysis calculations are automatically adjusted to reflect the length of test, the amount of moisture in the sample, temperature, time from the end of test, and the amount of radiation measured. If your test kit was used prior to the Use By date, ALL the testing protocols and instructions were carefully followed, and the data recorded properly on the test packet, then it is reasonable to assume this is an accurate assessment of the average level of the radon this sample was exposed to during the test period.

INTERPRETING YOUR TEST RESULT

This radon test was provided to you by BLUE RIDGE NET PUBLISHING INC / 540-777-1895. The US EPA action level for indoor radon is 4.0 pCi/L. Test results in this range (0.5 pCi/L or less) are, for all practical purposes, equivalent to the radon levels found in fresh air. However, if you make any structural changes or start to use a lower level of the building more frequently you should test again.

You may be able to obtain additional information about radon related subjects by calling the "Radon Fix-It Line" at 800-644-6999 Monday thru Friday between NOON and 8 pm EST.

Health Risks

The primary health risk from long-term exposure to radon is lung cancer. The risk of developing a lung cancer from radon exposure depends both on how much radon is present and how long you are exposed to radon. The higher the radon level or the longer the time of exposure, even if the levels are relatively low, the greater the risk. EPA has set an Action Level for radon at 4 pCi/L; however radon concentrations less than 4 pCi/L still pose some health risks. The Indoor Radon Abatement Act set a goal for indoor radon concentrations to equal the amount of radon found outdoors, which is estimated to be ~ 0.4 pCi/L.

Conducting Follow-up Measurements

USEPA protocol describes two general types of radon measurements: short-term tests conducted from 48 hours up to 90 days, and long-term tests that last from 91 to 365 days. Your first test (initial/screening) should be a short-term 'worst-case' screening to see if there is a potential for high exposure to radon. Screening tests should be conducted under closed-building conditions, in the lowest lived-in area in the house, because the highest concentrations of radon will usually be found in a room closest to the underlying soil. Tests made under these conditions are less likely to miss a house with a potential for high concentrations. On the other hand, if the results of worst-case screening tests are very low, there is a high probability that the average annual concentrations in the house are also low.

(Continued on Back)

Most states have a radon office to assist citizens with general questions about radon and radon reduction techniques. Many states maintain a list of licensed or certified radon testing and mitigation professionals. You can visit www.state-radon.info to find the list of state radon contacts, as well as links to additional radon resources in your area.

01/18/19 ACTIVATED CHARCOAL RADON TEST #8773372

Radon Test Result: < 0.3 ±0.2 pCi/L

Test Started 01/09/19 at 8:00 am
Test Ended 01/15/19 at 9:00 am
Closed house conditions maintained during test.

Location 1st Floor



PRESCHOOL ROOM 2
345 1ST ST
GERVAIS, OR 97026-

INTERPRETING YOUR TEST RESULT

This radon test was provided to you by BLUE RIDGE NET PUBLISHING INC / 540-777-1895. The US EPA action level for indoor radon is 4.0 pCi/L. Test results in this range (0.5 pCi/L or less) are, for all practical purposes, equivalent to the radon levels found in fresh air. However, if you make any structural changes or start to use a lower level of the building more frequently you should test again.

You may be able to obtain additional information about radon related subjects by calling the "Radon Fix-It Line" at 800-644-6999 Monday thru Friday between NOON and 8 pm EST.



Air Chek, Inc.
PO Box 2000
Naples, NC 28760

www.radon.com

Your Test Result

This result has been rounded to one-tenth (0.1) of a pCi/L (picocurie per liter). This test result reflects the amount of radon measured in this sample AFTER it arrived at our laboratory. All analysis calculations are automatically adjusted to reflect the length of test, the amount of moisture in the sample, temperature, time from the end of test, and the amount of radiation measured. If your test kit was used prior to the Use By date, ALL the testing protocols and instructions were carefully followed, and the data recorded properly on the test packet, then it is reasonable to assume this is an accurate assessment of the average level of the radon this sample was exposed to during the test period.

Health Risks

The primary health risk from long-term exposure to radon is lung cancer. The risk of developing a lung cancer from radon exposure depends both on how much radon is present and how long you are exposed to radon. The higher the radon level or the longer the time of exposure, even if the levels are relatively low, the greater the risk. EPA has set an Action Level for radon at 4 pCi/L; however radon concentrations less than 4 pCi/L still pose some health risks. The Indoor Radon Abatement Act set a goal for indoor radon concentrations to equal the amount of radon found outdoors, which is estimated to be ~ 0.4 pCi/L.

Conducting Follow-up Measurements

USEPA protocol describes two general types of radon measurements: short-term tests conducted from 48 hours up to 90 days, and long-term tests that last from 91 to 365 days. Your first test (initial/screening) should be a short-term 'worst-case' screening to see if there is a potential for high exposure to radon. Screening tests should be conducted under closed-building conditions, in the lowest lived-in area in the house, because the highest concentrations of radon will usually be found in a room closest to the underlying soil. Tests made under these conditions are less likely to miss a house with a potential for high concentrations. On the other hand, if the results of worst-case screening tests are very low, there is a high probability that the average annual concentrations in the house are also low.

(Continued on Back)

Most states have a radon office to assist citizens with general questions about radon and radon reduction techniques. Many states maintain a list of licensed or certified radon testing and mitigation professionals. You can visit www.state-radon.info to find the list of state radon contacts, as well as links to additional radon resources in your area.

Attention: P3161 / WES PRITT

Kit #: 8659778 Result: < 0.3 pCi/l

Location: 1st Floor 1

Gervais School

150 Douglas Ave

Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 12:00 pm

Started : 2018-03-19 at 1:00 pm

Ended : 2018-03-22 at 10:00 am

Hours/MST% : 69 hours 4.5% 70°F

Kit #: 8659780 Result: < 0.3 pCi/l

Location: 1st Floor

Gervais School

150 Douglas Ave

Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 11:00 am

Started : 2018-03-19 at 1:00 pm

Ended : 2018-03-22 at 10:00 am

Hours/MST% : 69 hours 6.0% 40°F

Kit #: 8659781 Result: < 0.3 pCi/l

Location: 1st Floor 4

Gervais School

150 Douglas Ave

Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 11:00 am

Started : 2018-03-19 at 1:00 pm

Ended : 2018-03-22 at 10:00 am

Hours/MST% : 69 hours 4.5% 40°F

Kit #: 8659782 Result: < 0.3 pCi/l

Location: 5

Gervais School

150 Douglas Ave

Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 12:00 pm

Started : 2018-03-19 at 1:00 pm

Ended : 2018-03-22 at 10:00 am

Hours/MST% : 69 hours 5.3% 50°F

Kit #: 8659783 Result: 0.8 ± 0.6 pCi/l

Location: 1st Floor 5a

Gervais School

150 Douglas Ave

Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 12:00 pm

Started : 2018-03-19 at 1:00 pm

Ended : 2018-03-22 at 10:00 am

Hours/MST% : 69 hours 6.8% 50°F

Kit #: 8659784 Result: < 0.3 pCi/l

Location: 1st Floor 6

Gervais School

150 Douglas Ave

Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 11:00 am

Started : 2018-03-19 at 1:00 pm

Ended : 2018-03-22 at 10:00 am

Hours/MST% : 69 hours 4.5% 50°F

Attention: P3161 / WES PRITT

Kit #: 8659785 Result: ????

Location: 1st Floor
Gervais School
150 Douglas Ave
Gervais, OR 97026-8913

Analysis Note : DNI

Analyzed : 2018-03-31 at 11:00 am
Started : 2018-03-19 at 1:00 pm
Ended : 2018-03-20 at 10:00 am
Hours/MST% : 21 hours 5.3% 50°F

Kit #: 8659786 Result: < 0.3 pCi/l

Location: 1st Floor 8
Gervais School
150 Douglas Ave
Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 12:00 pm
Started : 2018-03-19 at 1:00 pm
Ended : 2018-03-22 at 1:00 pm
Hours/MST% : 72 hours 5.3% 50°F

Kit #: 8659787 Result: ????

Location: 1st Floor 9
Gervais School
150 Douglas Ave
Gervais, OR 97026-8913

Analysis Note : UDI

Analyzed : 2018-04-02 at 3:00 pm
Started : 2018-03-19 at 1:00 pm
Ended : 2018-03-22 at 10:00 am
Hours/MST% : 69 hours 6.8% 50°F

Kit #: 8659788 Result: < 0.3 pCi/l

Location: 1st Floor 10
Gervais School
150 Douglas Ave
Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 11:00 am
Started : 2018-03-19 at 1:00 pm
Ended : 2018-03-22 at 10:00 am
Hours/MST% : 69 hours 6.1% 50°F

Kit #: 8659789 Result: 1.4 ± 0.6 pCi/l

Location: 11
Gervais School
150 Douglas Ave
Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 12:00 pm
Started : 2018-03-19 at 1:00 pm
Ended : 2018-03-22 at 10:00 am
Hours/MST% : 69 hours 5.3% 50°F

Kit #: 8659790 Result: ????

Location: 1st Floor 12
Gervais School
150 Douglas Ave
Gervais, OR 97026-8913

Analysis Note : UDI

Analyzed : 2018-03-31 at 11:00 am
Started : 2018-03-19 at 1:00 pm
Ended : 2018-03-22 at 10:00 am
Hours/MST% : 69 hours 6.0% 50°F

Attention: P3161 / WES PRITT

Kit #: 8659791 Result: < 0.3 pCi/l

Location: 1st Floor 16

Gervais School

150 Douglas Ave

Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 11:00 am

Started : 2018-03-19 at 2:00 pm

Ended : 2018-03-22 at 10:00 am

Hours/MST% : 68 hours 4.5% 50°F

Kit #: 8659792 Result: < 0.3 pCi/l

Location: 17

Gervais School

150 Douglas Ave

Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 11:00 am

Started : 2018-03-19 at 2:00 pm

Ended : 2018-03-22 at 10:00 am

Hours/MST% : 68 hours 5.2% 70°F

Kit #: 8659794 Result: < 0.3 pCi/l

Location: 18

Gervais School

150 Douglas Ave

Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 11:00 am

Started : 2018-03-19 at 2:00 pm

Ended : 2018-03-22 at 10:00 am

Hours/MST% : 68 hours 5.2% 70°F

Kit #: 8659795 Result: < 0.3 pCi/l

Location: 19

Gervais School

150 Douglas Ave

Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 1:00 pm

Started : 2018-03-19 at 2:00 pm

Ended : 2018-03-22 at 10:00 am

Hours/MST% : 68 hours 4.5% 50°F

Kit #: 8659796 Result: < 0.3 pCi/l

Location: 1st Floor 20

Gervais School

150 Douglas Ave

Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 11:00 am

Started : 2018-03-19 at 2:00 pm

Ended : 2018-03-22 at 10:00 am

Hours/MST% : 68 hours 5.2% 50°F

Kit #: 8659797 Result: < 0.3 pCi/l

Location: 1st Floor 21

Gervais School

150 Douglas Ave

Gervais, OR 97026-8913

Analysis Note : D

Analyzed : 2018-03-31 at 11:00 am

Started : 2018-03-19 at 2:00 pm

Ended : 2018-03-22 at 10:00 am

Hours/MST% : 68 hours 6.8% 50°F
